



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 631 779 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **94108921.1**

(51) Int. Cl.⁶: **A61K 31/22**

(22) Date of filing: **10.06.94**

(30) Priority: **22.06.93 IT RM930399**

(43) Date of publication of application:
04.01.95 Bulletin 95/01

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC
NL PT SE**

(71) Applicant: **AVANTGARDE S.p.A.**
4 Via Treviso
I-00040 Pomezia(Rome) (IT)

(72) Inventor: **Cavazza, Claudio**
Piazza Campitelli 2
I-00186 Roma (IT)
Inventor: **Cavazza, Paolo**
Viale dell'Umanesimo, 178
I-00144 Roma (IT)

(74) Representative: **Fassi, Aldo**
c/o CON LOR S.r.l.,
Via Renato Fucini, 14
D-20133 Milano (IT)

(54) **Use of esters of L-carnitine and acyl L-carnitine with hydroxyacids for producing pharmaceutical compositions for treating dermatoses.**

(57) The use is disclosed of esters of L-carnitine and acyl L-carnitines with hydroxyacids for producing pharmaceutical compositions suitable to be topically applied for treating dermatoses such as ichthyosis and psoriasis.

EP 0 631 779 A1

The present invention relates to the use of esters of L-carnitine and acyl L-carnitine with hydroxyacids for producing pharmaceutical compositions which contain such esters as active ingredients, suitable to be topically applied for the treatment of dermatoses. Particularly preferred are the esters of the following hydroxyacids:

- 5 α - hydroxybutyric acid
- α - hydroxyisobutyric acid
- β - hydroxybutyric acid
- γ - hydroxybutyric acid
- α - hydroxyisocaproic acid
- 10 α - hydroxyisovalenic acid
- malic acid, and
- tartronic acid.

Preferably, the acyl group is C_1 - 5 alkanoyl, particularly acetyl, propionyl, butyryl, isobutyryl, valeryl and isovaleryl.

- 15 Encompassed by the compounds to be used according to the present invention are both the inner salts and the salts of the aforesaid esters with pharmacologically acceptable acids.

- Pharmaceutically acceptable salts of the compound of formula (I) include, in addition to the inner salts, all pharmaceutically acceptable salts which are prepared by the addition of acid to L-carnitine, and which do not give rise to undesirable toxic or collateral effects. The formation of pharmaceutically acceptable acid
- 20 addition salts is well known in pharmaceutical technology.

Non-limiting examples of suitable salts include the chloride, bromide, orotate, acid aspartate, acid citrate, acid phosphate, fumarate, acid fumarate, lactate, maleate, acid maleate, acid oxalate, acid sulfate, glucose phosphate, tartrate and acid tartrate salts.

- 25 The esters of L-carnitine and the aforesaid alkanoyl L-carnitine with β -hydroxybutyric acid and the pharmacologically acceptable salts thereof are known compounds.

For instance, EP 0443996 A1 discloses the activity of these esters in inhibiting neuronal degeneration (as it occurs e.g. in Alzheimer's dementia and Parkinson's disease) and liver proteolysis and in the treatment of coma.

- Also the esters of L-carnitine and the aforesaid alkanoyl L-carnitine with γ -hydroxybutyric acid and the
- 30 pharmacologically acceptable salts thereof are known compounds (see e.g. EP 429403 A2 and EP 442850 A1). These esters are endowed with the same pharmacological properties as the β -hydroxybutyric acid esters.

- On the other hand, the esters of L-carnitine and aforesaid alkanoyl L-carnitines with hydroxyacids other than β - and γ -hydroxybutyric acid are novel compounds. Their preparation can be carried out similarly to
- 35 that of the known esters which is disclosed in the aforesaid European patent applications with only slight modifications which, depending on the selected hydroxyacid, will be apparent to any average-skilled expert in organic synthesis.

The preparation of some of the esters suitable for the dermatologic use according to the present invention is hereinbelow described.

40

EXAMPLE 1

Preparation of the ester of L-carnitine with gamma-hydroxybutyric acid (ST 701).

- 45 **Step a:** Preparation of the benzyl ester of gamma-bromobutyric acid.

Gamma-bromobutyric acid (3.3 g; 0.02 moles) was suspended in benzyl alcohol (15 mL). The suspension was cooled to 0°C and thionyl chloride (8 mL; 0.01 moles) was slowly added dropwise thereto.

- The resulting mixture was kept at room temperature for 16 hours, then concentrated under vacuum for
- 50 removing the thionyl chloride and distilled for removing the benzyl alcohol. The distillation residue was shown to be the title compound.

TLC exane 6 - AcOEt4 $R_f = 0.8$

NMR CDCl_3 δ 7.2(5H,s,aromatic); 5.0(2H,s, CH_2 -benzyl) 3.3(2H,t, CH_2COO); 2.6-2.0(4H,m, BrCH_2CH_2)

- 55 **Step b:** Preparation of L-carnitine ester with benzyl gamma-bromobutyrate

Carnitine inner salt (0.8 g; 0.005 moles) was suspended in 10 mL anhydrous dimethyl formamide. Benzyl ester of gamma-bromobutyric acid (1.3 g; 0.005 moles) was added to the suspension. The resulting

reaction mixture was kept under stirring at 60 °C for 48 hours under a nitrogen stream and then distilled under vacuum till complete solvent removal; 1.3 g of residue were obtained which was shown to be the title compound.

5 TLC CHCl₃ 4.2-H₂O 1.1-Isopr OH 0.7 - CH₃COOH 1.1 MetOH 2.8 R_F = 0.8
 NMR D₂O δ 7.4(5H,s,aromatic); 5.2(2H,s,CH₂-benzyl); 4.6(1H,m,CHOH); 4.2(2H,m,O-CH₂); 3.6-
 (2H,m,N + CH₂); 3.3(9H,s,(CH₃)₃ N +); 3.0 (2H,d,CH-CH₂COO); 2.6(2H,m,CH₂ CH₂COO);
 2.0(2H,m,CH₂ CH₂CH₂).

Step c: Preparation of the ester of L-carnitine bromide with gamma-hydroxybutyric acid.

10

The compound of step b (1.3 g) was dissolved in 20 mL of a 1:1 H₂O : EtOH mixture. The resulting solution was hydrogenated in the presence of 150 mg 10% Pd/C at 3 atmospheres of hydrogen for 2 hours. The mixture was filtered and concentrated under vacuum. 1 g of the title compound was obtained.
 TLC as in step b R_F = 0.6

15

Step d: Preparation of the ester of L-carnitine chloride with gamma-hydroxybutyric acid (ST 701).

The compound of step c (1 g) was eluted on 30 mL of AMBERLITE IRA 402 strongly basic resin activated to Cl⁻ form. The eluate was lyophilized. A highly hygroscopic solid was obtained.

20 NMR (D₂O) δ 4.2(2H,t,-CH₂O-); 3.5(2H,d,-N + CH₂-); 3.2(9H,s,(CH₃)₃ N +); 2.0(2H,d,CH₂COO); 2.4-
 (2H,m,CH₂COOH); 2.0(2H,m,CH₂-CH₂COOH).

[α]_D²⁵ = -13.2° (C = 1, H₂O)

HPLC

Spherisorb column - SCX 5M

25 Eluant KH₂PO₄ 0.005 M - CH₃CN (35-65); pH = 4.2

Flow rate 1 mL/min

Detector UV 205 nm

ST 701 R_T = 7.8

Carnitine R_T = 10.02 0.5%

30

35


40

45

50

55

Ester of L-carnitine chloride with gamma-hydroxybutyric acid, synthesis scheme



EXAMPLE 2**Preparation of the ester of acetyl L-carnitine with γ -hydroxybutyric acid (ST 793)**

- 5 Step a: Preparation of the benzyl ester of γ -bromobutyric acid (ST 793).

γ -bromobutyric acid (3.3 g; 0.02 moles) was suspended in benzyl alcohol (15 mL). The suspension was cooled to 0°C and thionyl chloride (8 mL; 0.01 moles) was slowly added dropwise thereto. The resulting mixture was kept at room temperature for 16 hours, then concentrated under vacuum to remove the
10 unreacted thionyl chloride and distilled to remove the benzyl alcohol. The distillation residue was purified by silica gel chromatography using hexane-AcOEt 98:2 as eluant.

TLC hexane $R_F = 0.2$

NMR CDCl_3 δ 7.2(5H,s,aromatic); 5.0(2H,s, CH_2 -benzyl) 3.3(2H,t, CH_2COO); 2.6-2.0(4H,m, BrCH_2CH_2)

- 15 Step b: Preparation of the ester of acetyl L-carnitine with benzyl γ -bromobutyrate.

Acetyl L-carnitine inner salt (1.62 g; 0.008 moles) was suspended in 12 mL anhydrous dimethyl formamide. γ -bromobutyric acid benzyl ester (2.05 g; 0.008 moles) was added to the suspension.

The resulting reaction mixture was kept under stirring for 24 hours under a nitrogen stream.

- 20 Ethyl ether was then added till complete precipitation of a compound which was filtered off. 3.43 g of the title compound were thus obtained.

TLC CHCl_3 4:2- H_2O 1:1-Isopr OH 0:7- CH_3COOH 1:1

MeOH 2:8 $R_F = 0.8$

HPLC

- 25 Column μ Bondapack C18


eluant KH_2PO_4 0.05 M- CH_3CN 70:30

Flow rate 1 mL/min

R_t 12.9

NMR D_2O δ 7.4 (5H,s,aromatic);

30



- 35 5.2(2H,s, CH_2 -benzyl); 4.4-4.0(4H,m,N + CH_2OCH_2) 3.5(9H,s,(CH_3)₃N +); 3.2(2H,d, $\text{CH-CH}_2\text{COO}$); 2.3(2H,m, $\text{CH}_2\text{CH}_2\text{COO}$); 2.0(5H,m + , $\text{CH}_2\text{CH}_2\text{CH}_2\text{COCH}_3$)

Step c: Preparation of the ester of acetyl L-carnitine bromide with γ -hydroxybutyric acid.

- 40 The compound of the step b (1 g) was dissolved in 20 mL absolute ethanol. The resulting solution was hydrogenated in the presence of 100 mg 10% Pd7C at 3 atmospheres of hydrogen concentrated under vacuum. 0.75 g of the title compound were obtained. Yield 98%.


TLC as in step b $R_F = 0.7$

- 45 Step d: Preparation of the ester of acetyl L-carnitine with γ -hydroxybutyric acid inner salt.

The compound of step c (1 g) was eluted on 30 mL of a strongly basic resin AMBERLITE IRA 402 activated in HCO_3^- form. The eluate was lyophilized. A highly hygroscopic solid was obtained.

NMR (D_2O):

50



55

4.2(2H,t,- CH_2O); 3.7(2H,d,-N + CH_2 -); 3.2(9H,s,(CH_3)₃N +); 2.8(2H,d, CH_2COO); 2.3-2.0-
(5H,m + s, CH_2COOH + COCH_3); 1.8(2H,m, $\text{CH}_2\text{CH}_2\text{COOH}$)
 $[\alpha]_D^{25} = -18.0$ (C = 1, H_2O)

HPLC

Column spherisorb - SCX 5M

Eluant KH_2PO_4 0.005 M - CH_3CN (35-65); pH = 4.2

Flow-rate 1 mL/min

5 Detector UV 205 nm

Rt = 8.83

TLC as in step b $R_F = 0.5$ **EXAMPLE 3**

10

Preparation of the ester of isovaleryl L-carnitine chloride with β -hydroxybutyric acid (ST 687)Step a: Preparation of the benzyl ester of β -hydroxybutyric acid 1.

15 β -hydroxybutyric acid sodium salt (1.2 g; 0.01 moles) was suspended in benzyl bromide (6 mL; 0.05 moles) 18 crown-6 (0.264 g) dissolved in 7 mL acetonitrile was added to the mixture.

The resulting solution was partly concentrated under a nitrogen stream and then kept under stirring at 80 °C for 90 minutes. To the cooled solution a mixture hexane- H_2O was added. The separated and dried organic phase was concentrated and then distilled under vacuum for removing the excess benzyl bromide.

20 1.1 g of solid residue were obtained which was identified to be the title compound. Yield 56%.

TLC CHCl_3 9 - MeOH 1 $R_F = 0.8$

Gas chromatography: column HP1 25 m; inner diameter 0.32

mm; film thickness 0.33 μm

carrier (He) flow-rate : 1 mL/min.

25 Make up gas 40 mL/min

Splitting ration 40 mL/min

Injector 220 °C

Detector (Fid) 280 °C

Column temperature 120 °C, 15 °C/min 250 °C

30 Rt = 9.36 compound

Rt = 4.84 no benzyl bromide

NMR CDCl_3 δ 7.3(5H,s,benzyl); 5.2(2H,s, CH_2 -benzyl); 4.2(1H,m,CH); 2.8(1H,s,broadOH); 2.5-(2H,d, CH_2COO); 1.2 (3H,d, CH_3)

35 Step b: Preparation of the acid chloride of isovaleryl L-carnitine chloride 2.

Thionyl chloride (7.7 mL; 0.1 moles) was added to isovaleryl L-carnitine chloride (10 g; 0.035 moles). The resulting mixture was kept at room temperature for 4 hours, then concentrated under vacuum to remove the thionyl chloride excess. The residue was washed three times with anhydrous ethyl ether.

40 The raw reaction product thus obtained was used in the subsequent step without further purification.


Step c: Preparation of the ester of isovaleryl L-carnitine chloride with β -hydroxybutyric acid benzyl ester 3.

45 The acid chloride of isovaleryl L-carnitine chloride (0.035 moles) of step b was dissolved in anhydrous tetrahydrofuran (25 mL). To the resulting solution the β -hydroxybutyric acid benzyl ester (7 g; 0.035 moles) of step a was added.

The reaction mixture was kept at 25 °C under stirring over night. Ethyl ether was then added thereto till complete precipitation. The solid thus obtained was filtered off and washed with ethyl ester. 14 g of the title compound were obtained. Yield 89%.

50 NMR D_2O δ 5.7(5H,m,benzyl); 5.5(1H,m,-CH-); 5.2(1H,m,COOCH); 5.0(2H,s, CH_2 benz.) 3.8-(2H,m,NCH₂); 3.2(9H,s,(H₃)₃N⁺); 2.8-2.5(4H,dd, CH_2 -COOCHCH₂COO); 2.2(2H,d,OCOCH₂)

55



1.2(3H,d,CH-CH₃);

5 0.8 (6H, d, CH $\begin{array}{c} \text{CH}_3 \\ \diagup \\ \text{CH} \\ \diagdown \\ \text{CH}_3 \end{array}$) ;

10 Step d: Preparation of the ester of isovaleryl L-carnitine chloride with β -hydroxybutyric acid

The compound of step c (14 g; 0.031 moles) was dissolved in H₂O-ethanol 1:1 (100 mL) and hydrogenated in the presence of 1.5 g 10% Pd/C at 4 atmospheres for two hours.

The reaction mixture was filtered, the filtrate concentrated to dryness under vacuum and the residue
15 crystallized from acetone-ethyl ether. 10 g of a hygroscopic compound were obtained.

TLC chloroform 4.2 IsoprOH 0.7 MeOH 2.8 H₂O 1 AcOH 1.1 R_f=0.7 [α]_D²⁵ = -21° (C=1, H₂O)

NMRD₂O

20 δ 5.7 (1H, m, CH $\begin{array}{c} \text{CH}_3 \\ \diagup \\ \text{CH} \\ \diagdown \\ \text{CH}_3 \end{array}$) ;

25 5.3(1H,m,-COOCH-); 3.8(2H,m,N+CH₂) 3.2(9H,s,(CH₃)₃N+); 2.8(2H,d,CH₂-COO); 2.6-(2H,d,CH₂COOH); 2.2(2H,d,OCOCH₂);

30 1.8 (1H, m, CH $\begin{array}{c} \text{CH}_3 \\ \diagup \\ \text{CH} \\ \diagdown \\ \text{CH}_3 \end{array}$) ;

1.2(3H,d,CHCH₃);

35

0.8 (6H, d, CH $\begin{array}{c} \text{CH}_3 \\ \diagup \\ \text{CH} \\ \diagdown \\ \text{CH}_3 \end{array}$) ;

40

HPLC

Column μ Bondapack-C₁₈

Eluant KH₂ PO₄ 0.05M-CH₃CN (85-15)

Detector UV λ = 205 nm

45 Flow-rate 1ml/min

R_t = 14-16 (the diastereoisomers are shown)

50

Elementary Analysis for C ₁₅ H ₃₀ NO ₆ Cl			
	C	H	N
calc.	50.6	8.4	3.9
found	48.93	8.36	3.49

55 The dermatoses which are suitably treated with the compositions of the present invention are in particular ichthyosis, psoriasis and those dermatoses which are induced by a defective keratinization, such as dandruff, acne and palmar and plantar hyperkeratosis.

Ichthyosis is a dermatosis characterized by generalized dryness, harshness and scaling of the skin. It may occur as a hereditary disease present at birth, or as a metabolic disorder associated with hypothyroidism or with the intake of drugs (such as butyrophens) inhibiting lipid synthesis, or as a paraneoplastic syndrome, manifestation of a tumor process involving internal organs.

5 Xeroderma, the mildest form of ichthyosis is neither congenital nor associated with systemic abnormalities.

It usually occurs on the lower legs of middle-aged or older patients, most often in cold weather and in patients who bathe frequently. There may be mild to moderate itching and an associated dermatitis due to detergents or other irritants.

10 The inherited ichthyoses, all characterized by excessive accumulation of scale on the skin surface, are classified according to clinical, genetic, and histologic criteria.

Known treatments of any form of ichthyosis comprise topically applying to the skin hydrating emollients. Furthermore, salicylic acid or vitamin A-containing ointments have been widely used.

15 A keratolytic agent particularly effective in removing the scale in ichthyosis vulgaris, lamellar ichthyosis and sex-linked ichthyosis contains 6% salicylic acid in a gel composed of propylene glycol, ethyl alcohol, hydroxypropylene cellulose and water.

Further known drugs for the treatment of this disorder include: 50% propylene glycol in water, hydrophilic petrolatum and water (in equal parts), and cold cream and an α -hydroxy acid (e.g. lactic and pyruvic acid) in various bases. In lamellar ichthyosis, 0.1% tretinoin (vitamin A acid; retinoic acid) cream has been utilized. None of these treatments has been found satisfactorily effective.

20 Hyperkeratosis is a thickening of the stratum corneum of the skin.

The treatment of choice is the topical application of drugs containing urea, propylene glycol or salicylic acid. Also in this case, none of the known treatment has proved to be satisfactorily effective.

It has now been found that the compounds of the present invention, when topically applied as solutions, 25 lotions, creams or ointments containing from 0.01% to 20%, preferably from 1% to 15% and most preferably from 2 to 10% by weight of at least one of the foregoing compounds, are potently effective in achieving complete remission of ichthyotic conditions in humans and in healing psoriasis and those disorders brought about by an altered keratinization, such as dandruff, acne and palmar and plantar hyperkeratosis.

30 It has also been found that, if the solutions, creams or ointments of the invention are applied regularly on a daily basis, within about two to three weeks the effected skin areas will return to normal conditions.

The compounds of formula (I) are prepared via a process whose steps are illustrated in the following reaction scheme, wherein R, R₁ and X have the previously defined meanings.

In order to prepare the compositions of this invention, at least one of the esters according to the 35 invention is preferably dissolved in water or ethanol initially. The solution thus prepared may be admixed in the conventional manner with commonly available ointment bases such as hydrophilic ointment (USP) or petrolatum (USP).

The water or ethanol used to dissolve the compounds according to this invention may range in concentration of from 1 to 30%, by volume, of the total composition. The compounds of this invention may 40 also be formulated in a solution or lotion form.

For instance, an ester according to the invention is dissolved directly in a mixture of water, ethanol and propylene glycol (40:40:20 by weight).

Some examples of the formulation are herein below described:

45 Formulation 1:5% solution

5 grams of an ester according to the invention were dissolved in 5 mL of water and the resulting solution admixed with 40 mL of ethanol and 20 mL of propylene glycol. Sufficient water was added to make 100 mL of formulation.

50

Formulation 2:5% ointment

5 grams of an ester according to the invention were admixed with 95 grams of USP grade hydrophilic ointment, until an uniform consistency resulted.

55

Claims

1. Use of esters of L-carnitine and acyl L-carnitines with hydroxyacids selected from the group consisting of α -hydroxybutyric acid, α -hydroxyisocaproic acid, α -hydroxyisovaleric acid, malic acid and tartronic acid, wherein the acyl group is C₁₋₅ alkanoyl selected from acetyl, propionyl, butyryl, isobutyryl, valeryl and isovaleryl and the pharmacologically compositions suitable to be topically applied for the treatment of dermatoses.
5
2. The use of claim 1, wherein the esters of L-carnitine and the acyl L-carnitines are in the form of inner salts.
10
3. A pharmaceutical composition suitable to be topically applied for treating dermatoses, which comprises an ester of claim 1 as active ingredient and a pharmacologically acceptable excipient therefor.
- 15 4. The composition of claim 3 for treating ichthyosis and psoriasis.
5. The composition of claim 3 for treating dermatoses brought about by defective keratinization.
6. The composition of claim 5 for treating dandruff, acne and palmar and plantar hyperkeratosis.
20
7. The composition of anyone of the claims 3-6 in the form of solution, lotion, ointment or cream.
8. The composition of claim 7 which comprises from 0.01% to 20%, preferably from 1% to 15%, most preferably from 2% to 10% by weight, of at least one of the compounds of claim 1.
25
9. Ester of L-carnitine and acyl L-carnitine with α -hydroxybutyric acid, α -hydroxyisobutyric acid, α -hydroxyisocaproic acid, α -hydroxyisovaleric acid, malic acid and tartronic acid, wherein the acyl group is C₁₋₅ alkanoyl selected from acetyl, propionyl, butyryl, isobutyryl, valeryl and isovaleryl.
- 30 10. The pharmacologically acceptable salt of the ester of claim 9.

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 94 10 8921

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.5)
P, Y	EP-A-0 596 838 (AVANTGARDE) * claim 1 * * page 2, line 45 - line 54 * ---	1-10	A61K31/22
Y	FR-A-2 654 618 (SEDERMA) * page 1, line 24 - line 28 * * claim 1 * * page 3, line 17 * ---	1-10	
Y	FR-A-2 654 619 (SEDERMA) * page 1, line 29 - page 2, line 22 * ---	1-10	
Y	US-A-4 839 159 (WINTER ET AL.) * column 2, line 28 - line 35 * * column 3, line 57 - line 63 * -----	1-10	
			TECHNICAL FIELDS SEARCHED (Int. CL.5)
			A61K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 3 October 1994	Examiner Gerli, P
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document			